

**SAMPLE FINAL EXAM ADVANCED MECHANICS,
January 2020, time: 2 hours**

Three problems (all items have a value of 10 points)

Remark 1 : Answers may be written in English or Dutch.

Remark 2: Write answers of each problem on separate sheets and add your name on them.

Problem 1

Three point masses m_1, m_2 and m_3 move in a three-dimensional space under influence of only gravitational forces that they exert on each other. The gravitational potential energy due to two point masses i and j is given by

$$V_{ij} = \frac{-Gm_i m_j}{|\mathbf{r}_i - \mathbf{r}_j|},$$

where G is the universal gravitational constant and \mathbf{r}_i the position vector of point mass m_i . Use as generalised coordinates the cartesian coordinates (x_i, y_i, z_i) of each mass point with respect to a fixed origin.

- a. Find the Hamiltonian function H for this system.
- b. Derive the Hamiltonian canonical equations for coordinate x_1 and its associated conjugate momentum $p_{1,x}$, where $p_{1,x}$ is the x -component of \mathbf{p}_1 .

(If you do not have the answer of item a, use

$$H = \alpha p_{1,x}^2 + \beta p_{1,x} + \frac{\gamma}{((x_1 - \hat{x})^2 + \rho^2)^{1/2}},$$

where $\alpha, \beta, \gamma, \hat{x}$ and ρ are constants.)


- c. How many of Hamilton's canonical equations of this system are independent? Explain your answer.
-

See next page for problem 2

Problem 2

A coin is steadily rolling on a perfectly rough surface (see figure). The coin is a thin circular disk with radius a , mass m , moment of inertia I with respect to axes in the plane of the coin and moment of inertia I_s along its symmetry axis.

The velocity of the centre of mass of the coin is \mathbf{v}_{cm} and its angular velocity is $\boldsymbol{\omega}$. The contact point between the coin and the surface is denoted by P and the origin O is at the centre of mass of the coin. Unit vector \mathbf{j}' is in the direction from P to O, unit vector \mathbf{k}' is along the symmetry axis of the coin and rolling occurs in the direction opposite to that of unit vector $\mathbf{i}' = \mathbf{j}' \times \mathbf{k}'$. Finally, unit vector \mathbf{k} points in the vertical direction and \mathbf{g} is gravity.



- a. The condition of perfect rolling means that the velocity in contact point P is zero. Use this to show that

$$\mathbf{v}_{cm} = -\mathbf{i}'a\omega_{z'} + \mathbf{k}'a\omega_{x'}$$

$$\text{where } \omega_{x'} = \boldsymbol{\omega} \cdot \mathbf{i}' \text{ and } \omega_{z'} = \boldsymbol{\omega} \cdot \mathbf{k}'.$$

- b. The angular velocity components are given by

$$\omega_{x'} = \dot{\theta}, \quad \omega_{y'} = \dot{\phi} \sin \theta, \quad \omega_{z'} = \dot{\psi} + \dot{\phi} \cos \theta,$$

with θ , ϕ and ψ the Eulerian angles. The meaning of θ is given in the figure.

Give the definition of angles ϕ and ψ and make a figure in which you sketch ϕ and ψ .

- c. Use the Lagrange formalism to show that the equations for the rolling coin read

$$(I + ma^2)\ddot{\theta} = I\dot{\phi}^2 \sin \theta \cos \theta - (I_s + ma^2)S\dot{\phi} \sin \theta - mga \cos \theta,$$

$$\frac{d}{dt} \left[I\dot{\phi} \sin^2 \theta + (I_s + ma^2)S \cos \theta \right] = 0,$$

$$\frac{dS}{dt} = 0,$$

$$\text{with } S = \dot{\psi} + \dot{\phi} \sin \theta.$$

- d. Note that $\theta = \pi/2$ (upright rolling coin), $\phi = 0$ and $S = \text{constant}$ is a solution of the equations of motion in item c.


Under what condition(s) is this a stable solution?

Hint: substitute $\theta = (\pi/2) + \theta'$, $\phi = \phi'$, with $\theta' \ll 1$ and $\phi' \ll 1$, in the equations of motion and maintain only terms that are linear in θ and in ϕ' .

See next page for problem 3

Problem 3

A light elastic spring of stiffness K is clamped at its upper end and supports a particle of mass m at its lower end. A second spring of stiffness K is fastened to the particle and, in turn, supports a particle of mass $2m$ at its lower end. Note: the system in its equilibrium configuration is subject only to gravitational force.



- Find the normal frequencies of the system for vertical oscillations about the equilibrium configuration.
- Find the normal coordinates.
If you have no answer of item a, describe the method to find these coordinates.
- Determine the general solution for $x_1(t), x_2(t)$.
If you have no answer to item b, describe the method to find this solution.

END

Equation sheet Advanced Mechanics for final exam (version 2019/2020)

A1. Goniometric relations:

$$\begin{aligned} \cos(2\alpha) &= \cos^2 \alpha - \sin^2 \alpha, & \cos(\alpha \pm \beta) &= \cos \alpha \cos \beta \mp \sin \alpha \sin \beta \\ \sin(2\alpha) &= 2 \sin \alpha \cos \alpha, & \sin(\alpha \pm \beta) &= \sin \alpha \cos \beta \pm \cos \alpha \sin \beta \end{aligned}$$

A2. Spherical coordinates r, θ, ϕ :

$$\begin{aligned} x &= r \sin \theta \cos \phi, & y &= r \sin \theta \sin \phi, & z &= r \cos \theta \\ dx dy dz &= r^2 \sin \theta dr d\theta d\phi \\ \mathbf{v} &= \mathbf{e}_r \dot{r} + \mathbf{e}_\theta r \dot{\theta} + \mathbf{e}_\phi r \dot{\phi} \sin \theta \\ \mathbf{a} &= \mathbf{e}_r (\ddot{r} - r \dot{\phi}^2 \sin^2 \theta - r \dot{\theta}^2) + \mathbf{e}_\theta (r \ddot{\theta} + 2\dot{r} \dot{\theta} - r \dot{\phi}^2 \sin \theta \cos \theta) \\ &\quad + \mathbf{e}_\phi (r \ddot{\phi} \sin \theta + 2\dot{r} \dot{\phi} \sin \theta + 2r \dot{\theta} \dot{\phi} \cos \theta) \end{aligned}$$

A3. Cylindrical coordinates R, ϕ, z :

$$\begin{aligned} x &= R \cos \phi, & y &= R \sin \phi, & z &= z \\ dx dy dz &= R dR d\phi dz \\ \mathbf{v} &= \mathbf{e}_R \dot{R} + \mathbf{e}_\phi R \dot{\phi} + \mathbf{e}_z \dot{z} \\ \mathbf{a} &= \mathbf{e}_R (\ddot{R} - R \dot{\phi}^2) + \mathbf{e}_\phi (2\dot{R} \dot{\phi} + R \ddot{\phi}) + \mathbf{e}_z \ddot{z} \end{aligned}$$

A4. $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B})$

A5. $(\mathbf{A} \times \mathbf{B}) \cdot \mathbf{C} = (\mathbf{B} \times \mathbf{C}) \cdot \mathbf{A} = (\mathbf{C} \times \mathbf{A}) \cdot \mathbf{B}$

A6. $\left(\frac{d\mathbf{Q}}{dt}\right)_{fixed} = \left(\frac{d\mathbf{Q}}{dt}\right)_{rot} + \boldsymbol{\omega} \times \mathbf{Q}$

B1. Noninertial reference frames:

$$\begin{aligned} \mathbf{v} &= \mathbf{v}' + \boldsymbol{\omega} \times \mathbf{r}' + \mathbf{V}_0 \\ \mathbf{a} &= \mathbf{a}' + \dot{\boldsymbol{\omega}} \times \mathbf{r}' + 2\boldsymbol{\omega} \times \mathbf{v}' + \boldsymbol{\omega} \times (\boldsymbol{\omega} \times \mathbf{r}') + \mathbf{A}_0 \end{aligned}$$

C1. Systems of particles:

$$\sum_i \mathbf{F}_i = \frac{d\mathbf{p}}{dt}, \quad \frac{d\mathbf{L}}{dt} = \mathbf{N}$$

C2. Angular momentum vector: $\mathbf{L} = \mathbf{r}_{cm} \times m\mathbf{v}_{cm} + \sum_i \bar{\mathbf{r}}_i \times m_i \bar{\mathbf{v}}_i$
where $\bar{\mathbf{r}}_i = \mathbf{r}_i - \mathbf{r}_{cm}$, $\bar{\mathbf{v}}_i = \mathbf{v}_i - \mathbf{v}_{cm}$

C3. Equations of motion for 2-particle system with central force:

$$\mu \frac{d^2 \mathbf{R}}{dt^2} = f(R) \frac{\mathbf{R}}{R}$$

with $\mu = m_1 m_2 / (m_1 + m_2)$ the reduced mass, \mathbf{R} relative position vector.

C4. Motion with variable mass:

$$\mathbf{F}_{ext} = m\dot{\mathbf{v}} - \mathbf{V}\dot{m}$$

with \mathbf{V} velocity of Δm relative to m .

D1. Moment of inertia tensor:

$$\mathbf{I} = \sum_i m_i (\mathbf{r}_i \cdot \mathbf{r}_i) \mathbf{1} - \sum_i m_i \mathbf{r}_i \mathbf{r}_i$$

D2. Moment of inertia about an arbitrary axis: $I = \tilde{\mathbf{n}} \mathbf{I} \mathbf{n} = mk^2$

D3. Formulation for sliding friction: $F_P = \mu_k F_N$

D4. Impulse and rotational impulse: $\mathbf{P} = \int \mathbf{F} dt = m\Delta \mathbf{v}_{cm}$, $\int N dt = Pl$
with l the distance between line of action and the fixed rotation axis.

E1. Transformation rule components of a real cartesian tensor, rank p , dimension N :

$$T'_{i_1 i_2 \dots i_p} = \alpha_{i_1 j_1} \alpha_{i_2 j_2} \dots \alpha_{i_p j_p} T_{j_1 j_2 \dots j_p}$$

F1. Euler equations: $N_1 = I_1 \dot{\omega}_1 + \omega_2 \omega_3 (I_3 - I_2)$
(other equations follow by cyclic permutation of indices)

G1. Lagrange's equations (first kind):

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_i} \right) = \frac{\partial L}{\partial q_i} + \lambda_k \frac{\partial f_k}{\partial q_i}$$

with $f_k(q_1, q_2, \dots, q_n, t) = 0$ constraints.

G2. Hamilton's variational principle:

$$\delta \int_{t_1}^{t_2} L dt = 0$$

G3. Hamiltonian function:

$$H = p_i \dot{q}_i - L$$

G4. Hamilton's canonical equations:

$$\dot{p}_i = -\frac{\partial H}{\partial q_i}, \quad \dot{q}_i = \frac{\partial H}{\partial p_i}$$
