

## Inleiding Topologie exam, January 28, 2015

**Note:** In all the questions below, please explain/prove your answers (e.g., in Exercise 1, part a., b., etc, please do not just say "yes" or "no". Also, in part e. or g. of the same exercise, do not just write down the final result, but also explain how you found it- e.g. by explaining your reasoning using pictures). There are three questions marked with a "\*"; they are probably more difficult than the rest. The final mark for the exam is the number of points that you collect, except for the case in which you collect more than 10 points, when the final mark will be 10 (in total, the exercises below are worth 11.5 points).

**Exercise 1.** Prove that, for any Hausdorff space  $(X, \mathcal{T})$ , any finite subset  $F \subset X$  is closed in  $X$ .  
(1 point)

**Exercise 2.** Consider the family  $\mathcal{B}$  of subsets of  $\mathbb{R}^2$  consisting of all the subsets of type  $(a, b) \times (a, b)$  with  $a < b$  real numbers:

$$\mathcal{B} = \{(a, b) \times (a, b) : a, b \in \mathbb{R}, a < b\}.$$

Let  $\mathcal{T}$  be the smallest topology on  $\mathbb{R}^2$  containing  $\mathcal{B}$ . We also consider

$$A = [0, 1] \times [0, 2] \subset \mathbb{R}^2.$$

- Is  $(\mathbb{R}^2, \mathcal{T})$  second countable? (0.5 points)
- Is  $(\mathbb{R}^2, \mathcal{T})$  Hausdorff? (0.5 points)
- Is the identity map  $\text{Id} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  continuous as a map from  $(\mathbb{R}^2, \mathcal{T})$  to  $(\mathbb{R}^2, \mathcal{T}_{\text{Eucl}})$ ? But as a map from  $(\mathbb{R}^2, \mathcal{T}_{\text{Eucl}})$  to  $(\mathbb{R}^2, \mathcal{T})$ ? (0.5 points)
- Is  $A$ , with the topology induced from  $\mathcal{T}$ , connected? Is it compact? (0.5 points)
- Let  $x = (0, 2) \in \mathbb{R}^2$  (the point of coordinates 0 and 2). Compute the closure of  $\{x\}$  in  $(\mathbb{R}^2, \mathcal{T})$ . (0.5 points)
- Show that the sequence  $(x_n)_{n \geq 1}$  given by

$$x_n = (\sin^2(n), \cos^4(n + 2015)) \in \mathbb{R}^2$$

is convergent in  $(\mathbb{R}^2, \mathcal{T})$  and has more than one limit. (0.5 points)

- Compute the interior and the closure of  $A$  in  $(\mathbb{R}^2, \mathcal{T})$ . (1 point)

h\* Show that any continuous map  $f : (\mathbb{R}^2, \mathcal{T}) \rightarrow \mathbb{R}$  must be constant. (1 point)

(total:  $6 \times 0.5 + 2 \times 1 = 5$  points)

**Exercise 3.** Consider the group of integers modulo 2,  $\mathbb{Z}_2 = \{\hat{0}, \hat{1}\}$ . Define the following action of  $\mathbb{Z}_2$  on the closed unit disk  $D^2 = \{z \in \mathbb{C} : |z| \leq 1\}$ :

$$\hat{0} \cdot z = z, \quad \hat{1} \cdot z = -z \quad (\text{for all } z \in D^2).$$

Prove that the resulting quotient  $D^2/\mathbb{Z}_2$  is homeomorphic to  $D^2$  (provide a complete argument; a "proof" based only on pictures is not enough to get the entire 1 point for this exercise).

**(1 point)**

**Exercise 4.** Let  $X$  be compact Hausdorff space and let  $C(X)$  be the space of real-valued continuous functions on  $X$ , endowed with the topology induced by the *sup*-metric.

Prove that if  $\mathcal{A} \subset C(X)$  is a dense subset of  $C(X)$ , then  $\mathcal{A}$  must be point-separating.

**(1 point)**

**Exercise 5.** For any algebra  $A$  over  $\mathbb{R}$  and any ideal  $I$  of  $A$ , we define

$$X(A, I) := \{\chi \in X_A : \chi(x) = 0 \quad \forall x \in I\} \subset X_A$$

and we endow it with the topology induced from the topology on the spectrum  $X_A$  of  $A$ . Show that:

- a. For any algebra  $A$  and any ideal  $I$  of  $A$ ,  $X(A, I)$  is closed in  $X_A$ . *(0.5 points)*
- b. Applied to  $A = \mathbb{R}[x, y]$  (polynomials in two variables) and  $I$  the ideal consisting of polynomials that are divisible by  $x^2 + y^2 - 1$ ,  $X(A, I)$  is homeomorphic to  $S^1$ . *(0.5 points)*
- c\*. Assume now that  $X$  is a compact Hausdorff space,  $Y \subset X$  and set

$$A = C(X), \quad I := \{f \in C(X) : f(y) = 0 \quad \forall y \in Y\}.$$

Show that  $X(A, I)$  is homeomorphic to the closure  $\bar{Y}$  of  $Y$  in  $X$  (where the last space is endowed with the topology induced from  $X$ ). *(1 point)*

**(total:  $2 \times 0.5 + 1 = 2$  points)**

**Exercise 6.** For each natural number  $n$  we consider a space  $X_n$  that is obtained by removing  $n$  distinct points from  $\mathbb{R}^2$ . We consider the 1-point compactification  $X_n^+$  and we denote by  $\infty_n \in X_n^+$  the point at infinity (so that  $X_n^+ = X_n \cup \{\infty_n\}$ ). Show that

- a.  $X_n^+$  can be embedded in  $\mathbb{R}^3$  (here you do not have to write down explicit formulas for the embedding, but please explain your reasoning using pictures and mention what result(s) you use in order to reach the final conclusion). *(0.5 points)*
- b\*. If  $X_n$  and  $X_m$  are homeomorphic, then  $n = m$ . *(1 point)*

**(total:  $0.5 + 1 = 1.5$  points)**