

Maat en Integratie A (WISB312) 21 april 2004

Question 1

Let $\phi : [A, B] \rightarrow [a, b]$ be a strictly increasing surjective continuous function. Suppose $\psi : [a, b] \rightarrow \mathbb{R}$ is non-decreasing, and $f : [a, b] \rightarrow \mathbb{R}$ a bounded ψ -Riemann integrable function. Define α and g on $[A, B]$ by

$$\alpha(y) = \psi(\phi(y)), g(y) = f(\phi(y))$$

Show that g is α -Riemann integrable and

$$\int_A^B g d\alpha = \int_a^b f d\psi$$

Question 2

Let $\{c_n\}$ be a sequence satisfying $c_n \geq 0$ for all $n \geq 1$, and $\sum_{n=1}^{\infty} c_n < \infty$. Let $\{s_n\}$ be a sequence of distinct points in (a, b) . Define a function ψ on $[a, b]$ by $\psi(x) = \sum_{n=1}^{\infty} c_n \mathbf{1}_{(s_n, b]}(x)$, where $\mathbf{1}_{(s_n, b]}$ is the indicator function of the interval $(s_n, b]$. Prove that any continuous function f on $[a, b]$ is ψ -Riemann integrable, and

$$\int_a^b f(x) d\psi(x) = \sum_{n=1}^{\infty} c_n f(s_n)$$

Question 3

Let $\Gamma \subseteq \mathbb{R}^n$. Recall that the inner Lebesgue measure of Γ is defined by

$$|\Gamma|_i = \inf\{|\mathcal{K}|_e : \mathcal{K} \subseteq \Gamma, \mathcal{K} \text{ compact}\}$$

Prove the following:

- (a) Γ is Lebesgue measurable if and only if $|\Gamma|_e = |\Gamma|_i$.
- (b) Γ is Lebesgue measurable if and only if $|\mathcal{A}|_e = |\Gamma \cap \mathcal{A}|_e + |\Gamma^c \cap \mathcal{A}|_e$ for all $\mathcal{A} \subseteq \mathbb{R}^n$.
- (c) If $\mathcal{A} \subseteq \Gamma$, and Γ is Lebesgue measurable, then $|\mathcal{A}|_e + |\Gamma \setminus \mathcal{A}|_i = |\Gamma|_i$

Question 4

Let E be a set and \mathcal{A} an algebra over E . Let $\mu : \mathcal{A} \rightarrow [0, 1]$ be a function satisfying

- (I) $\mu(E) = 1 = 1 - \mu(\emptyset)$,
- (II) if $A_1, A_2, \dots, \in \mathcal{A}$ are pairwise disjoint and $\bigcup_{n=1}^{\infty} A_n \in \mathcal{A}$ then

$$\mu\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \mu(A_n)$$

- (a) Show that if $\{A_n\}$ and $\{B_n\}$ are increasing sequences in \mathcal{A} such that $\bigcup_{n=1}^{\infty} A_n \subseteq \bigcup_{n=1}^{\infty} B_n$, then $\lim_{n \rightarrow \infty} \mu(A_n) \leq \lim_{n \rightarrow \infty} \mu(B_n)$
- (b) Let \mathcal{G} be the collection of all subsets G of E such that there exists an increasing sequence $\{A_n\}$ in \mathcal{A} with $G = \bigcup_{n=1}^{\infty} A_n$. Define $\bar{\mu}$ on \mathcal{G} by

$$\bar{\mu}(G) = \lim_{n \rightarrow \infty} \mu(A_n)$$

Where $\{A_n\}$ is an increasing sequence in \mathcal{A} such that $G = \bigcup_{n=1}^{\infty} A_n$. Show the following.

(i) $\bar{\mu}$ is well defined.

(ii) If $G_1, G_2 \in \mathcal{G}$, then $G_1 \cup G_2, G_1 \cap G_2 \in \mathcal{G}$ and

$$\bar{\mu}(G_1 \cup G_2) + \bar{\mu}(G_1 \cap G_2) = \bar{\mu}(G_1) + \bar{\mu}(G_2)$$

(iii) If $G_n \in \mathcal{G}$ and $G_1 \subseteq G_2 \subseteq \dots$, then $\bigcup_{n=1}^{\infty} G_n \in \mathcal{G}$ and

$$\bar{\mu}\left(\bigcup_{n=1}^{\infty} G_n\right) = \lim_{n \rightarrow \infty} \bar{\mu}(G_n)$$

(c) Define μ^* on $\mathcal{P}(E)$ (powerset of E) by

$$\mu^*(A) = \inf\{\bar{\mu}(G) : A \subseteq G, G \in \mathcal{G}\}$$

(i) Show that $\mu^*(A) = \bar{\mu}(G)$ for all $G \in \mathcal{G}$ and

$$\mu^*(A \cup B) + \mu^*(A \cap B) \leq \mu^*(A) + \mu^*(B)$$

for all subsets A, B of E . Conclude that $\mu^*(A) + \mu^*(A^c) \geq 1$ for all $A \subseteq E$.

(ii) Show that if $C_1 \subseteq C_2 \subseteq \dots$ are subsets of E and $C = \bigcup_{n=1}^{\infty} C_n$, then $\mu^*(C) = \lim_{n \rightarrow \infty} \mu^*(C_n)$.

(iii) Let $\mathcal{H} = \{B \subseteq E : \mu^*(B) + \mu^*(B^c) = 1\}$. Show that \mathcal{H} is a σ -algebra over E , and μ^* is a measure on \mathcal{H} .

(iv) Show that $\sigma(E; \mathcal{A}) \subseteq \mathcal{H}$. Conclude that the restriction of μ^* to $\sigma(E; \mathcal{A})$ is a measure extending μ , i.e. $\mu^*(A) = \mu(A)$ for all $A \in \mathcal{A}$.

Question 5

Let $\bar{\mathcal{B}}_{\mathbb{R}^N}$ be the Lebesgue σ -algebra over \mathbb{R}^N , $\mathcal{B}_{\mathbb{R}^N}$ the Borel σ -algebra over \mathbb{R}^N , and $\mathcal{B}_{\bar{\mathbb{R}}}$ the Borel σ -algebra over $\bar{\mathbb{R}} = [-\infty, \infty]$. Denote by $\lambda_{\mathbb{R}^N}$ the Lebesgue measure on $\bar{\mathcal{B}}_{\mathbb{R}^N}$. Let $f : \mathbb{R}^N \rightarrow [-\infty, \infty]$ be a Lebesgue measurable function (i.e. $f^{-1}(A) \in \bar{\mathcal{B}}_{\mathbb{R}^N}$ for all $A \in \mathcal{B}_{\bar{\mathbb{R}}}$). Show that there exists a function $g : \mathbb{R}^N \rightarrow [-\infty, \infty]$ which is Borel measurable (i.e. $g^{-1}(A) \in \mathcal{B}_{\mathbb{R}^N}$ for all $A \in \mathcal{B}_{\bar{\mathbb{R}}}$) such that

$$\lambda_{\mathbb{R}^N}(\{x \in \mathbb{R}^N : f(x) \neq g(x)\}) = 0.$$

Question 6

Let (E, \mathcal{B}, μ) be a measure space, and $f : E \rightarrow [0, \infty]$ be a measurable simple function such that $\int_E f d\mu < \infty$. Show that for every $\epsilon > 0$ there exists a $\delta > 0$ such that if $A \in \mathcal{B}$ with $\mu(A) < \delta$ then $\int_A f d\mu < \epsilon$.