



**Measure and Integration Final Exam**

**Due date: June 30, 2004**

**You must work on this exam individually. It is not allowed to discuss this exam with your fellow student.**

1. Let  $\nu$  be a  $\sigma$ -finite measure on  $(E, \mathcal{B})$ , and suppose  $E = \bigcup_{n=1}^{\infty} E_n$ , where  $\{E_n\}$  is a collection of pairwise disjoint measurable sets such that  $\nu(E_n) < \infty$  for all  $n \geq 1$ . Define  $\mu$  on  $\mathcal{B}$  by  $\mu(\Gamma) = \sum_{n=1}^{\infty} 2^{-n} \nu(\Gamma \cap E_n) / (\nu(E_n) + 1)$ .

- (a) Prove that  $\mu$  is a finite measure on  $(E, \mathcal{B})$ .  
(b) Show that for any  $\Gamma \in \mathcal{B}$ ,  $\nu(\Gamma) = 0$  **if and only if**  $\mu(\Gamma) = 0$ .  
(c) Find explicitly two positive measurable functions  $f$  and  $g$  such that

$$\mu(A) = \int_A f d\nu \text{ and } \nu(A) = \int_A g d\mu$$

for all  $A \in \mathcal{B}$ .

2. Suppose that  $\mu_i, \nu_i$  are finite measures on  $(E, \mathcal{B})$  with  $\mu_i \ll \nu_i$ ,  $i = 1, 2$ . Let  $\nu = \nu_1 \times \nu_2$  and  $\mu = \mu_1 \times \mu_2$ .

- (a) Show that  $\mu \ll \nu$ .  
(b) Prove that  $\frac{d\mu}{d\nu}(x, y) = \frac{d\mu_1}{d\nu_1}(x) \cdot \frac{d\mu_2}{d\nu_2}(y)$   $\nu$  a.e.

3. Let  $(E, \mathcal{B}, \mu)$  be a measure space.

- (a) Suppose  $f, g \in L^1(\mu)$  are such that  $\int_A f d\mu \leq \int_A g d\mu$  for all  $A \in \mathcal{B}$ . Show that  $f \leq g$   $\mu$  a.e.  
(b) Show that  $\mu$  is  $\sigma$ -finite **if and only if** there exists a **strictly** positive measurable function  $f \in L^1(\mu)$ .

4. Let  $(E, \mathcal{B}, \mu)$  be a measure space, and  $\{f_n\} \subseteq L^1(\mu)$ ,  $f \in L^1(\mu)$  be such that (i)  $f_n, f \geq 0$  for  $n \geq 1$ , (ii)  $\int_E f_n d\mu = \int_E f d\mu < \infty$  for  $n \geq 1$ , and (iii)  $f_n \rightarrow f$   $\mu$  a.e.

- (a) Show that  $\lim_{n \rightarrow \infty} \int_E (f - f_n)^+ d\mu = 0$ .  
(b) Prove that  $\lim_{n \rightarrow \infty} \sup_{A \in \mathcal{B}} \left| \int_A f_n d\mu - \int_A f d\mu \right| = 0$ .

5. Consider the measure space  $(\mathbb{R}, \mathcal{B}(\mathbb{R}), \lambda)$ , where  $\mathcal{B}(\mathbb{R})$  is the Borel  $\sigma$ -algebra, and  $\lambda$  Lebesgue measure.

(a) Let  $\mu$  be a probability measure on  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$  and  $f : \mathbb{R} \rightarrow [0, 1)$  a measurable function such that  $\mu\left(f^{-1}\left(\left[\frac{k}{2^n}, \frac{k+1}{2^n}\right)\right)\right) = \frac{1}{2^n}$  for  $n \geq 1$  and  $k = 0, \dots, 2^n - 1$ . Show that  $f \in L^2(\mu)$ , and determine the value of  $\|f\|_{L^2(\mu)}$ .

(b) Show that  $\lim_{n \rightarrow \infty} \int_0^n \left(1 - \frac{x}{n}\right)^n e^{x/2} d\lambda(x) = 2$ .

(c) Let  $f : \mathbb{R} \rightarrow \overline{\mathbb{R}}$  be measurable, and suppose  $\int_{\mathbb{R}} f(x) d\lambda(x)$  exists. Show that for all  $a \in \mathbb{R}$ , one has

$$\int_{\mathbb{R}} f(x-a) d\lambda(x) = \int_{\mathbb{R}} f(x) d\lambda(x).$$

(d) Let  $k, g \in L^1(\lambda)$ . Define  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ , and  $h : \mathbb{R} \rightarrow \overline{\mathbb{R}}$  by

$$F(x, y) = k(x-y)g(y) \text{ and } h(x) = \int_{\mathbb{R}} F(x, y) d\lambda(y).$$

(i) Show that  $F$  is measurable.

(ii) Show that  $\lambda(|h| = \infty) = 0$  and  $\int_{\mathbb{R}} |h| d\lambda \leq \left(\int_{\mathbb{R}} |k| d\lambda\right) \left(\int_{\mathbb{R}} |g| d\lambda\right)$ .

6. Consider the measure space  $([a, b], \mathcal{B}, \lambda)$ , where  $\mathcal{B}$  is the Borel  $\sigma$ -algebra on  $[a, b]$ , and  $\lambda$  is the restriction of the Lebesgue measure on  $[a, b]$ . Let  $f : [a, b] \rightarrow \mathbb{R}$  be a bounded Riemann integrable function. Show that the Riemann integral of  $f$  on  $[a, b]$  is equal to the Lebesgue integral of  $f$  on  $[a, b]$ , i.e.

$$(R) \int_a^b f(x) dx = \int_{[a,b]} f d\lambda.$$